

**Washington State University**  
***Institutional Animal Care and Use Committee***

**Guideline #9: Blood Collection**

**Approval Date: 08/09/2021 (Replacing version 08/2018)**

---

**A. Purpose**

This document is intended to provide guidance regarding safe volumes and common routes for blood collection from laboratory animals.

---

**B. Guideline**

The volume of blood collected for research purposes from most mammals is generally not a problem. However, repeated blood sample collection in mice, rats, hamsters, guinea pigs, small cats, and birds can be problematic because of their small body size. In order to prevent anemia, electrolyte imbalance, hypovolemic shock or other complications, the following guidelines should be followed:

1. The acceptable quantity and frequency of blood sampling is determined by the circulating blood volume and the red blood cell (RBC) turnover rate.
2. For optimal health, blood draws should be limited to the lower end of the range. Maximum blood volumes should be taken only from healthy animals.
3. The approximate circulating blood volume of an animal is 55-70 ml/kg. A blood volume estimate for a single species may not reflect differences among individual breeds or variations due to age, size, or illness. In older or obese animals, circulating blood volume can be decreased as much as 15%.
4. Of the circulating blood volume, approximately 10% of the total volume can be safely removed every 2 to 4 weeks, 7.5% every 7 days, and 1% every 24 hours. The same total amount of blood can also be removed as multiple quantities over a 24-hour period. See **Table 1** for estimated blood volume for individual species. **Table 2** includes collection volumes for mice and rats of various weights. **Table 3** lists possible blood collection sites for multiple species.
5. With provision of replacement fluids, (equal volume of blood removed) (0.9% saline, Lactated Ringer's solution), up to 15% of circulating blood volume may be

collected at one time. This would require a 4-week recovery period before additional blood draws.

6. **When an increase of blood volume to be collected is requested for an IACUC approved protocol, anything over 10% every 2 weeks would need to go through a regular amendment process and would not qualify for a Veterinary Verification Consultation (VVC) according to WSU IACUC Policy #24 ([https://iacuc.wsu.edu/documents/2016/06/policy\\_24.pdf/](https://iacuc.wsu.edu/documents/2016/06/policy_24.pdf/)).**
7. Although blood *volume* is rapidly restored in an animal after blood collection, the rest periods described above are needed for blood *constituents* (e.g., red blood cells, platelets, clotting factors) to be regenerated by the body. Hemostasis after collection can be achieved by using a silver nitrate stick, Quick Stop powder or by applying a gauze sponge over the site with gentle pressure until bleeding stops.

**Table 1: Circulating Blood Volume & Examples of Maximal Survival Collection Calculation.**

Species	Circulating Blood Volume (mL/kg)	Average Adult Body Weight	Blood Volume	Maximum Survival Collection Volume (10% of circulating blood volume)
Mouse	77-80	25 g	2 mL	0.2 mL or 200 µL
Rat	50-70	300 g	18 mL	1.8 mL
Hamster	78	85-150 g	9.2 mL	0.9 mL
Gerbil	67	45-130 g	5.9 mL	0.59 mL
Guinea pig	67-92	700-1200 g	75 mL	7.5 mL
Rabbit	57-65	2.5 kg	143 mL	14 mL
Cat	47-60	4 kg	214 mL	21 mL
Dog (beagle)	79-90	12 kg	1 L	100 mL
Ferret	75	1 kg	75 mL	7.5 mL
Pig (large)	65	110 kg	7.15 L	715 mL
Sheep	60	60 kg	3.6 L	360 mL
Goat	70	45 kg	3.15 L	315 mL
Cattle	60	520 kg	31.2 L	3.12 L
Horse	75	400 kg	30 L	3 L
Chicken	60	2.5 kg	150 mL	15 mL

**Table 2: Approximate Blood Sample Volume Ranges and Safe Frequency of Collection for Mice and Rats.**

Body Weight (g)	Total Blood Volume (mL)	1% (mL)	7.5% (mL)	10% (mL)	15% (mL)*
<b>Mouse</b> based on mean 78 mL/kg (blood volume/body weight)					
20	1.56	0.02	0.12	0.16	0.23
25	1.95	0.02	0.15	0.20	0.29
30	2.34	0.02	0.18	0.23	0.35
35	2.73	0.03	0.20	0.27	0.41
40	3.12	0.03	0.23	0.31	0.47
<b>Rat</b> based on mean 60 mL/kg (blood volume/body weight)					
125	7.50	0.08	0.56	0.75	1.13
150	9.00	0.09	0.68	0.90	1.35
200	12.00	0.12	0.90	1.20	1.80
250	15.00	0.15	1.13	1.50	2.25
300	18.00	0.18	1.35	1.80	2.70
350	21.00	0.21	1.58	2.10	3.15
Frequency of Collection		Every 24 hours	Every 7 days	Every 2 weeks	Every 4 weeks

\*With provision of replacement fluids. See item 6 under Guideline.

**Table 3: Blood collection sites for multiple species.**

Species	Site
<b>Mouse</b>	Saphenous, tail veins or arteries, retro-orbital (requires anesthesia, less invasive methods preferred), submandibular and submental
<b>Rat</b>	Saphenous, tail veins or arteries, pedal and jugular veins
<b>Hamster</b>	Saphenous and jugular veins
<b>Guinea pig</b>	Saphenous, tarsal and jugular veins, vena cava
<b>Rabbit</b>	Ear vein or artery and jugular veins

<b>Ferret</b>	Jugular, cephalic and saphenous veins, cranial vena cava
<b>Cat</b>	Jugular, cephalic and saphenous veins
<b>Dog</b>	Saphenous, cephalic and jugular veins
<b>Pig</b>	Ear, saphenous, tail, cephalic, femoral and mammary veins, vena
<b>Sheep &amp; Goat</b>	Jugular and cephalic veins
<b>Cattle</b>	Jugular veins
<b>Horse</b>	Jugular veins
<b>Chicken</b>	Brachial and jugular veins

If you have any questions about this guideline, need training in one of the above methods, or need the information regarding a species not listed above, please call the Office of the Campus Veterinarian at 509-335-6246 or email at [or.ocv.alert@wsu.edu](mailto:or.ocv.alert@wsu.edu).

### C. References

---

1. Diehl K, et al. 2001, A Good Practice Guide to the Administration of Substances and Removal of Blood, Including Routes and Volumes, *J Appl Toxicol* 21:15-23.
2. Iwarsson K, Lindberg L, Waller T, 1994, Common non-surgical techniques and procedures. Chapter 16. Svensen P, Hau J (eds). In: *Handbook of Laboratory Animal Science*. Volume 1. CRC Press, Inc. Boca Raton, FL.
3. National Research Council, 2011. *Guide for the Care and Use of Laboratory Animals*. 8th Edition. The National Academies Press, Washington, DC